



CYTO

GUARANTEED MINIMUM ANALYSIS

6-Furfurylaminopurine (Kinetin).....	0.90%
Salicylic Acid.....	0.30%
Vitamin B1.....	0.30%
Vitamin C.....	0.30%

EZ-Gro Cyto contains biostimulants and plant growth regulators found in nature. It stimulates seedling and plant growth and may increase quality and yield. **EZ-Gro Cyto** helps to reduce transplant shock and increases root weight, and the number and size of flowers.

ATTENTION

Keep out of reach of children and animals
Due to low pH, may cause skin, eye and respiratory irritation

Wear protective clothing, gloves, eye and respiratory protection while handling product

Do not apply within 21 days of harvest | Avoid re-entry within 12 hours of foliar application. | Do not allow rinse water to contaminate streams, ponds and lakes as water life may be endangered. | For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. | Do not connect an irrigation system (including greenhouse systems) used for pesticide application to public water system. | If exposed, concerned, or feeling unwell contact a health care provider. | Store locked up in original container, in a well ventilated area. Keep container tightly closed. | Dispose of contents and container in accordance with local, regional, and national regulations.

LOT NUMBER:

EXPIRY DATE:

APPLICATION CHART:

CROP	APPLICATION USE ENOUGH WATER FOR COVERAGE
Barley, Oats, Rye, Spring or Winter Wheat, Canola, Flaxseed, Mustard, Safflower, Sunflower and Corn	3-5 leaf stage rate 110 ml/ha
Legumes (Soybean, Beans, Lentils, Chickpeas, Peas)	
Alfalfa, Clover, Hay	1st application 110 ml/ha at 3-5 leaf stage. 2nd application after cut.
Kale, Cabbage, Broccoli, Cauliflower, Brussel Sprouts	
Fruiting Vegetables (Tomatoes, Peppers, Eggplant)	Apply 110 ml/ha 14-21 days after emergence. 2nd application at 5-10% bloom.
Cucumbers, Watermelon, Squash	
Leafy Vegetables (Lettuce, Celery, Spinach, Swiss Chard)	Apply 110 ml/ha 14-21 days after emergence. 2nd application mid-season.
Sugar Beets	Apply 110 ml/ha 14-21 days after emergence.
Apples	1st application full pink 290 ml/ha. 2nd application petal fall 290 ml/ha. 3rd application 21 days after petal fall 290 ml/ha. 4th application 50 days after petal fall 290 ml/ha.
Stone Fruit (Peaches, Cherries, Plums, Apricots)	1st application pre-bloom 290 ml/ha. 2nd application petal fall 290 ml/ha. 3rd application 21 days after petal fall 290 ml/ha.
Grapes	1st application leaf out/pre-bloom 140 ml/ha. 2nd application petal fall 290 ml/ha.
Flowers & Bedding Species	1st application at transplant. 2nd application at 5-10% bloom. 0.45 ml/EZ-Gro K to 1 L water (110 ml/ha).
Turf & Sod Production	Apply 8.25 ml/100 m ² every 3-4 weeks foliarly to greens & tees.
Hydroponics	Add 1 ml/10 L water to reservoir feed weekly through transplant, grow and bloom.
Cannabis	Add 1 ml/4 L water in the last 2 weeks of Grow and first week of Bloom, followed by 3ml/4 L water in weeks 2-3 of Bloom, and 6ml/4L water in weeks 4 and 5 of Bloom.