



FERTIACTYL® GZ 13-0-5

GUARANTEED ANALYSIS

F 1550

• Total Nitrogen (N)... 13% 13 % Urea Nitrogen

· Soluble Potash (K2O) ... DERIVED FROM: Urea, Potassium Hydroxide

NON PLANT FOOD INGREDIENTS:

Humic acid Fulvic acid Seaweed extract

PLEASE READ CAREFULLY BEFORE USING - KEEP OUT OF REACH OF CHILDREN - RECIRCULATE BEFORE USE

DIRECTIONS FOR USE:

- Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Add FERTIACTYL® GZ first in a half full clean sprayer and agitate while mixing. Keep agitating until the end of the application. Dilute into at least 20 Gal/Acre.

• FERTIACTYL® GZ is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants.

CROP	N° OF APP.	APPLICATION STAGES	APP. RATE (pts/Acre)	ADVICE
Corn, Soybean & Alfalfa	1-4	At planting	2-4	Broadcast app.
			1.5 - 4	In row app.
		Early growth	2 - 4	Broadcast app.
			1.5 - 4	Banded app.
Cotton	2	At planting		Apply in furrows.
			2-4	• 4 pts/Acre gives best results.
		With herbicide application		• It can be added to liquid starter fert. with glyphosate foliar app. Total should be 4 pts/Acre.
General Vegetables	1-3	At planting	0.25 - 1% transplant solution	Must have mechanical or by pass tank agitation in transplant water tank.
		At transplant to		May be mixed with compatible fertilizers (not calcium nitrate) or injected in
		7 days after planting	2-4	clean water (drip irrigation). If not available, use it to water plants (same rate),
		With 2nd foliar spray	1	wetting leaves until the solution reaches the soil.
		(7 - 14 days after 1st application)		
Wheat and Small Grains	1-2	Planting to FS 3	1-3	Tank mixes well with most pesticides and liquid fertilizers.
				• Enhances tillering.

COMPATIBILITY AND STORAGE:

- FERTIACTYL® GZ is compatible with most liquid fertilizers and a wide range of agrochemicals. Do not mix with low pH products or calcium nitrate. In all cases, test the compatibility prior to treatment. Manufacturer's responsibility is limited to the guarantee of product's composition, as per label in certificate packaging. Do not store under 32°F (0°C).

Volume: 264 Gallons (1000 liters) - Net weight: 2667 lb (1210 kg)

TIMAC AGRO USA, INC. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of TIMAC AGRO USA, INC. In no case shall TIMAC AGRO USA, INC. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. TIMAC AGRO USA, INC. makes no warranties of merchantability or fitness for the standard product of the sa particular purpose nor any other express or implied warranty except as stated above.

MANUFACTURED BY TIMAC AGRO USA -P.O. Box 888 - Route 724 & I-176 Reading, PA 19607 -0888 - USA

